arXiv:astro-ph/0610004v2 23 Oct 2006

Submitted to the *Astrophysical Journal Letters*: Revised

Discovery of a Candidate Central Compact Object in the Galactic Nonthermal SNR G330.2+1.0

Sangwook Park[1], Koji Mori[2], Oleg Kargaltsev[1], Patrick O. Slane[3], John P. Hughes[4], David N. Burrows[1], Gordon P. Garmire[1], and George G. Pavlov[1]

ABSTRACT

We report on the discovery of a pointlike source (CXOU J160103.1−513353) at the center of a Galactic supernova remnant (SNR) G330.2+1.0 with *Chandra X-Ray Observatory*. The X-ray spectrum fits a black-body (BB) model with $kT \sim 0.49$ keV, implying a small emission region of $R \sim 0.4$ km at the distance of 5 kpc. The estimated X-ray luminosity is $L_X \sim 1 \times 10^{33}$ ergs s^{-1} in the $1 - 10$ keV band. A power law model may also fit the observed spectrum, but the fit results in a very large photon index, $\Gamma \sim 5$. We find no counterparts at other wavelengths. The X-ray emission was steady over the ∼13 hr observation period, showing no variability. While we find marginal evidence for X-ray pulsations ($P \approx 7.5$ s), the presence of a pulsar at the position of this object is not conclusive with the current data, requiring an independent confirmation. These results are generally consistent with an interpretation of this object as a Central Compact Object associated with SNR G330.2+1.0.

Subject headings: stars: neutron — X-ray: stars — ISM: individual (SNR G330.2+1.0) — supernova remnants

1. INTRODUCTION

Since the discovery of nonthermal X-ray emission from the shell of supernova (SN) 1006 (Koyama et al. 1995), the nonthermal X-ray emitting blast wave shock fronts of supernova remnants (SNRs) have been considered as prime candidate sites for the cosmic-ray acceleration. Thanks to the high-resolution, high-sensitivity detectors on board *Chandra* and *XMM-Newton* observatories, several SNRs now show evidence for particle acceleration sites near the shock fronts. While

[1]Department of Astronomy and Astrophysics, Pennsylvania State University, 525 Davey Laboratory, University Park, PA. 16802; park@astro.psu.edu

[2]Department of Applied Physics, University of Miyazaki, 1-1 Gakuen Kibana-dai Nishi, Miyazaki, 889-2192, Japan

[3]Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA. 02138

[4]Department of Physics and Astronomy, Rutgers University, 136 Frelinghuysen Road, Piscataway, NJ. 08854-8019

the nonthermal X-ray emission is typically from localized regions near the shock front in young historical SNRs (e.g., Cas A, Tycho, SN 1006), nonthermal X-rays in some relatively "older" ($\tau \gtrsim$ a few 10^3 yr) SNRs dominate the entire face of the remnants (G347.3−0.5 and G266.2−1.2, Koyama et al. 1997; Slane et al. 2001). Each of the latter also contains a Central Compact Object (CCO) (Slane et al. 1999; Pavlov et al. 2001), a peculiar manifestation of isolated neutron stars (NSs) found in near the center of only a handful of SNRs (e.g., Pavlov et al. 2004).

Recently, using the archival *ASCA* data, Torii et al. (2006, T06 hereafter) discovered that the X-ray emission from a Galactic shell-type radio SNR G330.2+1.0 (Caswell et al. 1983) is dominated by nonthermal continuum over the entire SNR with enhancements toward the boundary shell. The X-ray continuum was described by a power law (PL) model (photon index $\Gamma \sim 2.8$), indicating the X-ray synchrotron emission from the shock-accelerated electrons. Based on the distance of $\sim$5 − 10 kpc (McClure-Griffiths et al. 2001), a Sedov age of $\tau \gtrsim 3000$ yr was estimated, assuming the typical explosion energy ($E_0 \sim 10^{51}$ ergs) and ambient density ($n_0 \sim 1$ cm^{-3}) (T06). G330.2+1.0 appears to resemble G347.3−0.5 and G266.2−1.2, but, unlike those two SNRs, no central point source was found for G330.2+1.0 in the *ASCA* data (T06).

We observed G330.2+1.0 with *Chandra* in order to perform a detailed study of the SNR such as the spatially-resolved spectroscopy of the nonthermal X-ray emitting shell, which could not be performed with the low-resolution detectors of *ASCA*. The other major goal was the search for a NS at the center of the SNR, motivated by the fact that the other two nonthermal X-ray-dominated SNRs, G347.3−0.5 and G266.2−1.2, contain a CCO. We report here the discovery of a pointlike source at the center of G330.2+1.0, which shows characteristics similar to the known CCOs. In this *Letter*, we present the results from the analysis of X-ray emission from this new candidate CCO. The results from the analysis of the SNR are presented elsewhere (Park et al., in preparation).

2. OBSERVATIONS & DATA REDUCTION

SNR G330.2+1.0 was observed with the Advanced CCD Imaging Spectrometer (ACIS) on board *Chandra X-Ray Observatory* in May 22, 2006 as part of the Guaranteed Time Observations program. G330.0+1.0 has an $\sim$10′ angular size (Caswell et al. 1983, T06). It is heavily absorbed and dominated by nonthermal continuum in the hard X-ray band ($E = 1 - 10$ keV). We thus used the ACIS-I array to utilize its large field of view (FOV) of $\sim$17′ × 17′ and the relatively large collecting area in the hard band. The pointing was roughly toward the geometrical center of the SNR. No strong variability was found in the background light curve. We corrected for the spatial and spectral degradation of the ACIS data caused by radiation damage, known as the charge transfer inefficiency (CTI; Townsley et al. 2000), with the methods developed by Townsley et al. (2002a), before further standard data screening by status, grade, and energy selections. "Flaring" pixels were removed, and the standard grades (02346) were selected. After the data reduction, the effective exposure was 49.5 ks. The overall SNR spectrum is hard, and there are few source photons below $E \sim 1$ keV. The central pointlike source, as discussed below, also shows a spectrum with

nearly all source photons detected at $E > 1$ keV. We thus extracted photons in the $1 - 7$ keV band for the data analysis. (The upper bound of $E = 7$ keV was chosen to avoid instrumental lines and the higher background rates at $E > 7$ keV.)

3. Data Analysis

The broadband image of SNR G330.2+1.0 is presented in Fig. 1. We detect some ∼20 pointlike sources within the SNR boundary (∼5′ radius). Among these sources, the one detected at the "center" of the SNR is remarkable. This source is the brightest (∼0.012 counts s^{-1}) among the sources detected within the SNR shell (all other sources are fainter by $\gtrsim$ an order of magnitude.) The source position is nearly at the geometrical center of the SNR: e.g., only ∼33″ southeast of the radio SNR center (Fig. 1) (Caswell et al. 1983). As we discuss below, this source appears to be a candidate NS associated with G330.2+1.0. Thus, we hereafter refer this object as the "NS candidate". We note that the NS candidate uniquely stands out in the central region of the SNR: e.g., within ∼1′ radius of the SNR center, we find only one other source at ∼45″ northeast of the NS candidate, which shows a soft spectrum and an optical counterpart and is thus most likely a foreground star. Comparisons of the NS candidate radial profile with the ACIS PSF at the source position show no evidence for an extended nebulosity, confirming the pointlike nature. The detected sky position of the source is RA(2000) = $16^h\ 01^m\ 3\overset{s}{.}14$, Dec(2000) = $-51^\circ\ 33'\ 53\overset{''}{.}6$. It is detected within ∼1′ off-axis on the ACIS-I2, and thus the positional uncertainties are ∼0″.3 (∼90% C. L.) (Alexander et al. 2003). We designate this source as CXOU J160103.1−513353.

We extracted the source spectrum (∼600 counts) from a circular region with a radius of 2″. The background spectrum was taken from a surrounding annular region (the inner radius = 4″ and the outer radius = 15″). The source spectrum was binned to contain a minimum of 15 counts per energy bin. For the spectral analysis of our CTI-corrected data, we utilized the response matrices appropriate for the spectral redistribution of the CCD, as generated by Townsley et al. (2002b). The X-ray spectrum is continuum-dominated with no clear line features (Fig. 2). A single blackbody (BB) model with $kT \sim 0.49$ keV can adequately describe the observed spectrum (Table 1). The high foreground column ($N_H \sim 2.5 \times 10^{22}$ cm^{-2}) is consistent with that for the SNR (T06, Park et al. in preparation). Alternatively, a PL model can fit the data, implying an extremely steep spectrum ($\Gamma \sim 5$) and a higher $N_H \sim 4.7 \times 10^{22}$ cm^{-2}. For completeness, we note that optically thin thermal plasma models (`mekal`) may also describe the spectrum with $kT \sim 0.95$ keV. The fit, however, requires an unrealistically low metal abundance with a 90% upper limit of ∼0.3 solar, which supports a continuum-dominated nature of the spectrum.

The NS candidate shows no significant evidence of variability on time scales of $\sim 1 - 7$ hr (Fig. 3). Although our *Chandra* observation was perfomed with the standard 3.24 s frame-time, which precludes the search for an ordinary pulsar-like periodicity ($P \lesssim 1$ s), we can attempt to search for a long periodicity of $P \sim$ several seconds, typical for Anomalous X-ray Pulsars (AXPs) (Mereghetti et al. 2002). The ACIS time resolution of 3.24 s and the total exposure of ∼50 ks

allow us to search for pulsations in the $4 \times 10^{-5} - 0.15$ Hz range. We use the arrival times of the 603 photons (of which ∼99.8% are expected to come from the source), recalculated to the solar system barycenter using the `axBary` tool. We used the Z^2_m test (Buccheri et al. 1983) to search for periodic pulsations. We calculated Z^2_m for $m = 1 - 4$ (where m is the number of harmonics included) at 1.5×10^5 equally spaced frequencies f in the $4 \times 10^{-5} - 0.15$ Hz range. This corresponds to oversampling by a factor of about 20, compared to the expected width of $T^{-1}_{\rm span} \approx 20$ μHz of the $Z^2_m(f)$ peaks, and guarantees that we miss no peaks. The most significant peak was found in Z^2_2 (= 29.5, corresponding to the ≈95% significance level) at $f = 0.133663\,{\rm Hz} \pm 5\,\mu{\rm Hz}$ ($P \approx 7.48$ s) (Fig. 4). If this corresponds to the actual period of the source, the pulsed fraction is ≈30%. However, we note that the statistical significance of the period is only marginal. If this period is simply due to the statistical fluctuation, we place a 3σ upper limit of ≈52% on the pulsed fraction, assuming sinusoidal pulse shape per period. Follow-up observations with higher time resolution and better photon statistics will be essential to verify the suggested periodicity or to search for a different period.

4. Discussion

The sky position at the center of SNR G330.2+1.0 and the high foreground column, which is consistent with that of the SNR, make the NS candidate association with the SNR highly plausible. There is no evidence for hour-scale variability that might be caused by orbital and/or stellar activities. We find no counterparts at other wavelengths. There is no radio (843 and 1415 MHz) enhancement at the SNR center (Caswell et al. 1983), and the nearest cataloged radio pulsar (PSR B1558−50) is ∼36′ away. The nearest optical source (USNO B1.0 #0384−0529508, B = 16.92, R = 15.08) is ∼3.″7 off the NS candidate. Considering the accuracy of both the X-ray and the optical positions ($\lesssim$0.″3), this source is unlikely to be the optical counterpart of the NS candidate. The optical upper limit (V ≈ 21) then implies a large X-ray-to-optical flux ratio of $f_{X(1-10\ {\rm keV})}/f_V > 9$, which supports the NS interpretation rather than a normal stellar or extragalactic object (Stocke et al. 1991; Agüeros et al. 2006). The high absorbing column toward G330.2+1.0 supports previously estimated distances of $d \sim 5 - 10$ kpc. Assuming that the NS candidate is associated with G330.2+1.0, we scale the distance to the NS candidate in units of 5 kpc.

The best-fit BB temperature of the NS candidate is higher than those expected from the standard cooling of the young NS's surface (Yakovlev & Pethick 2004), but it is consistent with the temperature observed in CCOs and/or AXPs (Mereghetti et al. 2002; Pavlov et al. 2002, 2004). The small size of the implied BB radiating region, $R_{BB} \sim 0.4\ d_5$ km (where d_5 is the distance scaled to 5 kpc), is similar to that of other CCOs (Pavlov et al. 2000,2002; Kargaltsev et al. 2002), probably indicating the X-ray emission from a hot polar cap of the NS. The X-ray luminosity $L_{X(1-10\ {\rm keV})} \sim 1 \times 10^{33}\ d_5^2$ ergs s^{-1} is also consistent with the object being a CCO (Pavlov et al. 2004 and references therein). The PL model implies a higher N_H than in the BB case and a very steep slope of the spectrum. These parameters resemble those of the CCOs in the other two non-

thermal X-ray SNRs G347.3−0.5 and G266.2−1.2 (Kargaltsev et al. 2002; Lazendic et al. 2003). The higher column associated with the PL fit could suggest a background AGN origin of the NS candidate, as perhaps suggested by the relatively high chance probability (∼0.2) of such a detection within the SNR boundary (∼5′ radius) based on the LogN-LogS relation in the Galactic plane (Ebisawa et al. 2001). However, the combination of the extremely steep X-ray spectrum ($\Gamma \sim 5$), the high f_X/f_O ratio, the absence of a radio counterpart, and the suggested possible pulsations strongly favor the case of a NS, making an AGN interpretation highly unlikely.

The NS candidate was not detected with *ASCA* (T06). The non-detection might be attributed to the low sensitivity (e.g., the GIS count rate is ∼30% of the ACIS-I at $E > 1$ keV) and the poor angular resolution (∼3′ FWHM) of *ASCA*. In fact, the ∼20 ks exposure of the archival *ASCA* data implied only ∼70 photons per GIS over the $\gtrsim$10 arcmin^2 source area. On the other hand, we serendipitously detected the NS candidate in archival *XMM-Newton* data (ObsID 0201500101). Assuming a BB model with $kT = 0.5$ keV and $N_H = 2.5 \times 10^{22}$ cm^{-2}, we estimate $L_X \sim 1.3 \times 10^{33}$ ergs s^{-1} for the *XMM-Newton* source[1], which is consistent with our *Chandra* results.

Based on the overall spectral/temporal properties and the position at the center of SNR G330.2+1.0, we propose that CXOU J160103.1−513353 is a candidate CCO associated with G330.2+1.0. The CCOs are believed to be a peculiar manifestation of NSs, whose nature is not fully understood. Considering only a handful of the known CCOs, the discovery of this new candidate CCO is important as it adds a new member to this class of NSs. The association with a newly-discovered nonthermal SNR G330.2+1.0 makes this new CCO candidate particularly interesting because each of the other two nonthermal X-ray-dominating SNRs also contains a CCO. Furthermore, the possible pulsations at $P \sim 7.5$ s (if confirmed), which is typical for AXPs, makes this object potentially even more intriguing, because only two of the known CCOs show pulsations, both with shorter periods (Zavlin et al. 2000; Gotthelf et al. 2005). If the pulsations with the long period are confirmed with follow-up observations, the NS candidate may considered to represent a "bridging" class between CCOs and AXPs.

This work was supported in part by SAO under *Chandra* grant SV4-74018. OK and GGP were supported by NASA grant NAG5-10865. KM was partially supported by the Grant in-Aid for Young Scientists (B) of the MEXT (No. 18740108).

REFERENCES

Agüeros, M. A. et al. 2006, AJ, 131, 1740

Alexander, D. M. et al. 2003, AJ, 126, 539

[1]The short exposure of ∼8 ks provided only limited photon statistics (∼50 counts per MOS) and did not allow further analysis. It was off the pn FOV due to the use of the small window mode.

Buccheri, R., et al. 1983, A&A 128, 245

Caswell, J. L., Haynes, R. F., Milne, D. K., & Wellington, K. J. 1983, MNRAS, 204, 915

Ebisawa, K., Maeda, Y., Kaneda, H., & Yamauchi, S. 2001, Science, 293, 1633

Gotthelf, E. V., Halpern, J. P., & Seward, F. D. 2005, ApJ, 627, 390

Kargaltsev, O., Pavlov, G. G., Sanwal, D., & Garmire, G. P. 2002, ApJ, 580, 1060

Koyama, K., Petre, R., Gotthelf, E. V., Hwang, U., Matsuura, M., Ozaki, M., & Holt, S. S. 1995, Nature, 378, 255

Koyama, K., Kinugasa, K., Matsuzaki, K., Nishiuchi, M., Sugizaki, M., Torii, K., Yamauchi, S., & Aschenbach, B. 1997, PASJ, 49, L7

Lazendic, J. S., Slane, P. O., Gaensler, B. M., Plucinsky, P. P., Hughes, J. P., Galloway, D. K., & Crawford, F. 2003, ApJ, 593, L27

McClure-Griffiths, N. M., Green, A. J., Dickey, J. M., Gaensler, B. M., Haynes, R. F., & Wieringa, M. H. 2001, ApJ, 551, 394

Mereghetti, S., Chiarlone, L., Israel, G. L., & Stella, L. 2002, Proc. of the 270. WE-Heraeus Seminar on Neutron Stars, Pulsars, and Supernova Remnants, MPE Report 278, eds. W. Becker, H. Lesch & J. Trümper (Garching bei München: MPE), p. 29

Pavlov, G. G., Zavlin, V. E., Aschenbach, B., Trümper, J., & Sanwal, D. 2000, ApJ, 531, L53

Pavlov, G. G., Sanwal, D., Kiziltan, B., & Garmire, G. P. 2001, ApJ, 559, L131

Pavlov, G. G., Sanwal, D., Garmire, G. P., & Zavlin, V. E. 2002, in Neutron Stars in Supernova Remnants, ASP Conf. Ser., Vol. 271, eds., P. O. Slane & B. M. Gaensler (San Francisco: ASP), p. 247

Pavlov, G. G., Sanwal, D., & Teter, M. 2004, in Young Neutron Stars and Their Environments, IAU Symposium 218, eds., F. Camilo & B. Gaensler (San Francisco: ASP), 239

Slane, P. O., Gaensler, B. M., Dame, T. M., Hughes, J. P., Plucinsky, P. P., & Green, A. 1999, ApJ, 525, 357

Slane, P. O., Hughes, J. P., Edgar, R. J., Plucinsky, P. P., Miyata, E., Tsunemi, H., & Aschenbach, B. 2001, ApJ, 548, 814

Stocke, J. T., Morris, S. L., Gioia, I. M., Maccacaro, T., Schild, R., Wolter, A., Fleming, T. A., & Henry, P. J. 1991, ApJS, 76, 813

Torii, K., Uchida, H., Hasuike, K., Tsunemi, H., Yamauchi, Y., & Shibata, S. 2006, PASJ, in press (astro-ph/0601569) (T06)

Townsley, L. K., Broos, P. S., Garmire, G. P., & Nousek, J. A. 2000, ApJ, 534, L139

Townsley, L. K., Broos, P. S., Chartas, G., Moskalenko, E., Nousek, J. A., & Pavlov, G. G. 2002a, Nucl. Instrum. Methods Phys. Res. A, 486, 716

Townsley, L. K., Broos, P. S., Nousek, J. A., & Garmire, G. P. 2002b, Nucl. Instrum. Methods Phys. Res. A, 486, 751

Yakovlev, D. G. & Pethick, C. J. 2004, ARA&A, 42, 169

Zavlin, V. E., Pavlov, G. G., Sanwal, D., & Trümper, J. 2000, ApJ, 540, L25

This preprint was prepared with the AAS LaTeX macros v5.2.

Table 1. Best-Fit Spectral Parameters of the NS Candidate.

Model	N_H 10^{22} cm^{-2}	Γ	kT keV	$f_{X(1-7\ \mathrm{keV})}$ [a] 10^{-13} ergs cm^{-2} s^{-1}	$L_{X(1-10\ \mathrm{keV})}$ 10^{33} d_5^2 ergs s^{-1}	R_{BB} [b] d_5 km	χ^2/ν
BB	$2.5^{+0.5}_{-0.4}$	-	$0.49^{+0.04}_{-0.04}$	1.17±0.10	1.0	$0.40^{+0.34}_{-0.17}$	28.4/33
PL	$4.7^{+0.7}_{-0.6}$	$5.0^{+0.6}_{-0.5}$	-	1.23±0.10	6.3	-	25.0/33

Note. — Errors are with 90% confidence.

[a]Obeserved X-ray flux.

[b]Black-body radius.



Fig. 1.— Broadband (1 − 7 keV) ACIS image of G330.2+1.0. Darker gray-scales correspond to higher intensities. For the purposes of display, the image has been binned with 4 pixels (∼2″) and then smoothed by a Gaussian with $\sigma = 5$ pixels. The NS candidate (CXOU J160103.1−513353) is marked with an arrow. The lower-right inset is the unbinned (0.″5 pixel) image of the NS candidate. The radio SNR center (Caswell et al. 1983) is marked with "+".



Fig. 2.— X-ray spectrum of the NS candidate as observed with the ACIS. The best-fit BB model is overlaid. The lower panel is the residuals from the best-fit BB model.



Fig. 3.— X-ray light curve of the NS candidate (filled circles). For comparison, the light curve from the southwestern filament of SNR G330.2+1.0 is presented with crosses. Each light curve has been binned with 2500 s.



Fig. 4.— Pulse profile of the NS candidate folded with the frequency of 0.133663 Hz ($P = 7.48$ s).